



Industry
Canada

Industrie
Canada

Canada

Français
Home

Contact Us
Site Map

Help
What's New

Search
About Us

Canada Site
Registration

Strategis Index:

A B C D E F G H I J K L M N O P Q R S T U V W X Y
Z

strategis.gc.ca

CIPO  OPIC



Canadian Patents Database

09/12/2002 - 13:04:44

(12) Patent Application:

(11) CA 2254060

(54) EFFERVESCENT BISPHOSPHONATE FORMULATION

(54) FORMULATION DE BISPHOSPHONATE EFFERVERSCENTE

[View or Download Images](#)

ABSTRACT:

An effervescent formulation of alendronate contains an acid source, a carbonate source, a binder, a lubricant and, optionally, flavoring agents, colorants and sweeteners.

CLAIMS: [Show all claims](#)

*** Note: Data on abstracts and claims is shown in the official language in which it was submitted.

(72) Inventors (Country):	KATDARE, Ashok V. (United States) KRAMER, Kenneth A. (United States) GARDNER, Colin R. (United States)
(73) Owners (Country):	MERCK & CO., INC. (United States)
(71) Applicants (Country):	MERCK & CO., INC. (United States)
(74) Agent :	OGILVY RENAULT
(45) Issued :	
(22) Filed :	May 13, 1997
(43) Laid Open :	Nov. 27, 1997
Examination requested :	May 8, 2002
(51) International Class (IPC) :	A61K 9/46 A61K 31/66

[Patent Cooperation Treaty \(PCT\)](#): Yes

(85) [National Entry](#):

Nov. 10, 1998

(86) <u>PCT Filing number:</u>	PCT/US1997/008055
(87) <u>International publication number:</u>	WO1997/044017

(30) Application priority data:

Application No.	Country	Date
60/017,881	United States	May 17, 1996
9613183.4	U.K.	June 24, 1996

Availability of licence:**N/A**Language of filing:**English**

View or Download Images :

- ☒ Cover Page Image
- ☐ Abstract Image
- ☐ Claims Image
- ☐ Disclosures Image
- ☐ Drawings Image
- ☐ Representative Drawing Image

[View the Image](#)[Download in Adobe PDF](#)[Top of Page](#)

Last Modified: 2001/06/13

[Important notices and disclaimer](#)
[Privacy Statement](#)

TITLE OF THE INVENTION
EFFERVESCENT BISPHOSPHONATE FORMULATION

CROSS-REFERENCE TO RELATED APPLICATIONS

5 Not Applicable

STATEMENT REGARDING FEDERALLY-SPONSORED R&D
Not Applicable

10 **REFERENCE TO MICROFICHE APPENDIX**
Not Applicable

FIELD OF THE INVENTION

15 This invention relates to pharmaceutical effervescent
formulations of bisphosphonates, especially alendronate, and in
particular, tablets and powders which effervesce when added to water.

BACKGROUND OF THE INVENTION

20 There are numerous bisphosphonates which have been
identified as having utility as pharmaceutical agents which inhibit bone
resorption. These include:

alendronate-- (4-amino-1-hydroxy-butylidene)bis-phosphonate;
cimadronate-- [(cycloheptylamino)methylene]bis-phosphonate;
clodronate-- (dichloromethylene)-bis-phosphonate;
25 EB-1053-- [1-hydroxy-3-(1-pyrrolidiny)-propylidene]bis-
phosphonate;
etidronate-- (1-hydroxyethylidene)-bis-phosphonate;
ibandronate-- [1-hydroxy-3-(methylpentylamino)propylidene]bis-
phosphonate;
30 neridronate-- (6-amino-1-hydroxyhexylidene)bis-phosphonate;
olpadronate-- [3-(dimethylamino)-1-hydroxy-propylidene]bis-
phosphonate;
pamidronate-- (3-amino-1-hydroxypropylidene)bis-phosphonate;
risedronate-- [1-hydroxy-2-(3-pyridinyl)-ethylidene]bis-
35 phosphonate;

tiludronate-- [[(4-chlorophenyl)thio]methylene]bis-phosphonate
YH 529-- [1-hydroxy-2-imidazo-(1,2a)pyridin-3-ylethylidene]bis-
phosphonate; and
zoledronate-- [1-hydroxy-2-(1H-imidazol-1-yl)ethylidene]bis-
5 phosphonate.

Alendronate has been approved by various regulatory agencies, including the Food and Drug Administration in the United States as an oral osteoporosis treatment in post menopausal women. The
10 currently marketed formulation is a tablet, and the patient is instructed to take the tablet with a full glass of water in the morning, at least a half hour prior to eating or drinking. However, certain side effects, including esophageal irritation and erosion have been reported if the tablet was not taken with enough water, or if the patient did not remain in an upright
15 position for approximately one-half hour after taking the medication.

It would be desirable to develop a formulation which allows a bisphosphonate to be taken orally, yet avoids problems associated with prolonged contact between the medication and the esophagus.

20 SUMMARY OF THE INVENTION

Not Applicable

BRIEF DESCRIPTION OF THE DRAWINGS

Not Applicable

25

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to an effervescent pharmaceutical formulation comprising, as an active ingredient, a bisphosphonate selected from the group consisting of: alendronate,
30 cimadronate, clodronate, EB-1053, etidronate, ibandronate, neridronate, olpadronate, pamidronate, risedronate, tiludronate, YH 529, zoledronate, pharmaceutically acceptable salts and esters of the foregoing, and mixtures of the foregoing;

an acid source selected from the group consisting of: citric acid, tartaric acid, malic acid, fumaric acid, adipic acid, succinic acid; an anhydride of said acids; an acid salt selected from the group consisting of sodium dihydrogen phosphate, disodium dihydrogen pyrophosphate and sodium acid sulfite; and mixtures of the acids, anhydrides and acid salts;

a carbonate source selected from the group consisting of: sodium bicarbonate, sodium carbonate, potassium bicarbonate, potassium carbonate, sodium sesquicarbonate, sodium glycine carbonate, and mixtures thereof;

a binder selected from the group consisting of a non-reducing sugar such as mannitol, lactose, dextrose or sorbitol; polyvinyl pyrrolidone, or, sodium chloride, sodium benzoate and sodium sulfate;

a lubricant selected from the group consisting of stearic acid salts, powdered sodium benzoate, L-leucine, sodium laurel sulfate and polyethylene glycol (PEG); and

optionally, one or more additional agents selected from the group consisting of flavoring agents (such as orange and cherry) colorants, and sweeteners (including xylitol, aspartame or acesulfame K).

The present invention is also directed to an effervescent pharmaceutical formulation comprising alendronate or a pharmaceutically acceptable salt thereof as an active ingredient and

an acid source selected from the group consisting of: citric acid, tartaric acid, malic acid, fumaric acid, adipic acid, succinic acid; an anhydride of said acids; an acid salt selected from the group consisting of sodium dihydrogen phosphate, disodium dihydrogen pyrophosphate and sodium acid sulfite; and mixtures of the acids, anhydrides and acid salts;

a carbonate source selected from the group consisting of: sodium bicarbonate, sodium carbonate, potassium bicarbonate, potassium carbonate, sodium sesquicarbonate, sodium glycine carbonate, and mixtures thereof;

wherein the acid source is present in at least an equivalent amount relative to the source of carbonate;

a binder selected from the group consisting of mannitol, sorbitol, polyvinyl pyrrolidone, lactose, sodium chloride sodium benzoate and sodium sulfate;

a lubricant selected from the group consist of stearic acid salts, powdered sodium benzoate, L-leucine, sodium laurel sulfate, powdered sodium bicarbonate and polyethylene glycol (PEG); and optionally, one or more additional agents selected from the group consisting of flavoring agents, particularly orange and cherry colorants, and sweeteners selected from the group consisting of xylitol, aspartame and acesulfame K.

The effervescent pharmaceutical formulation of the present invention may be either a tablet or a powder. To use the formulation, the tablet or powders are placed in an convenient amount of water to produce an effervescent liquid, and the patient drinks the effervescent liquid.

In one embodiment the formulation is a tablet, wherein in one class the total weight of the tablet ranges from about 100 to about 50,000 mg. In another embodiment, the tablet weight ranges from about 1500 to about 32,500 mg and more particularly from about 20,800 to about 30,150 mg.

As used throughout this specification and claims, the term "bisphosphonate" is intended to include the related bisphosphonic acids and salts, and various crystalline and amorphous forms. "Alendronate" includes the related bisphosphonic acid, and salt forms. It includes crystalline, hydrated crystalline, and amorphous forms of alendronate. It specifically includes alendronate sodium and alendronate monosodium trihydrate.

Methods for the preparation of bisphosphonates are well known in the art. Methods for the preparation of alendronate and alendronate sodium salt trihydrate are known. In particular, methods for the preparation of alendronate may be found in U.S. patents 4,922,007, 5,019,651 and 5,510,517, each of which is hereby incorporated by reference. The amount of active in the formulation based on a weight of alendronate will range from 1 to 80 mg, particularly 5-40 mg and more

particularly 5-15 mg. Exemplary amounts of alendronate are 5, 10 and 40 mg.

The formulations of this invention have many distinct advantages. First of all, the patients drinks an effervescent liquid, which
5 limits the amount of time in which the bisphosphonate is in contact with the esophageal tissue, thus minimizing the risk of irritation. Secondly, the bioavailability of at least some of the bishosphonates, including alendronate is increased as a consequence of an excipient acting as a sequestering agent. Thirdly, the alendronate formulations are particularly
10 advantageous in that alendronate is often prescribed to elderly patients who may experience difficulty in swallowing pills, but can more easily swallow a liquid formulation.

In preferred embodiments of the invention the acid source is chosen from acid sources which are also sequestering agents. This is an
15 important consideration because bisphosphonates, particularly alendronate, can be a potent sequestering agent of divalent cations, especially Ca^{2+} and Mg^{2+} . If either of these cations are present, the alendronate will sequester them, rendering the alendronate less bioavailable. Preferred acid sources which also act as a sequestering
20 agent include citric acid and tartaric acid, and mixtures thereof. The excess citric acid or tartaric acid in the formulation binds the ions in issue and prevents their complexation with alendronate.

The carbonate source should be chosen so that it does not contain divalent cations which could be sequestered by the
25 bisphosphonate. Suitable carbonate sources are sodium bicarbonate, sodium carbonate, potassium bicarbonate, potassium carbonate and sodium glycine carbonate. Preferred carbonate sources are sodium bicarbonate, sodium carbonate, and mixtures thereof.

In one aspect of this invention the acid source is present in
30 an amount equal or greater than the carbonate source, on a molecular equivalent basis. Thus, when citric acid is the acid source and sodium bicarbonate is the carbonate source, the mole ratio of citric acid/bicarbonate is at least 1:3. In a particular class of this aspect there is an excess of citric acid, as this not only helps to efficiently generate the

effervescence, but also acts to sequester any ions which might otherwise complex with alendronate and the excess also acts as a flavor enhancer. In an illustration of the excess acid source, the mole ratio of citric acid to bicarbonate ranges from 1:1 to 3:1. Where sodium carbonate is used as
5 the source of carbonate an equal equivalent with citric acid will require a mole ratio of 2 moles citric acid to 3 moles carbonate. Analogous ratios can be calculated for any source of acid and carbonate, and of course the carbonate source may be present as a mixture of bicarbonate and carbonate.

10 For patients who exhibit gastric irritation another aspect of this invention is to employ excess amounts of the carbonate source to provide an antacid effect to the formulation. Example 2 below demonstrates such a formulation.

The binder may be a non-reducing sugar, such as mannitol,
15 compressible lactose, dextrose or sorbitol. Alternatively, a polymer such as polyvinylpyrrolidone may be used. Other binders can include sodium chloride, sodium benzoate and sodium sulfate. The binder is usually present in approximately 3-10% of the tablet, and preferably about 5% (based on wt/wt), e.g. if the tablet weighs 1000 mg, the binder will be
20 from about 30-100 mg, more preferably about 50 mg. If, however, a direct compression process is used to make the tablets, then a higher amount of binder is employed, preferably about 300-400 mg.

The tablet should also contain a lubricant, which typically comprises about 0.5 to about 5% of the tablet on a wt/wt basis, and
25 preferably about 1%. Preferred lubricants include powdered sodium benzoate, micronized polyethylene glycol 6000 and polyethylene glycol 8000, sodium lauryl sulfate, powdered sodium bicarbonate and L-leucine.

For effervescent powder formulations, the composition of the powder is similar to that of the tablet, except that the lubricant is
30 generally present in a lesser amount (about less than one-half that in the tablet) and the binder should be chosen so that it is a dry binder. In preferred formulations, the powder is granulated.

Where desired a colorant, flavoring agent such as orange or cherry or sweetner from the group xylitol, aspartame, or acesulfame K

may be added to the formulation. Various methodologies for preparing the formulations of the current invention are illustrated in the examples below.

The following formulations and manufacturing procedures can be used for manufacture of effervescent tablets containing alendronate sodium.

It should be understood that one skilled in this art will recognize equivalent formulations which are intended to be included with the scope of this invention.

EXAMPLE 1

	Alendronate sodium (in mg alendronate acid)	5-10	10 mg
15	Citric acid, anhydrous (granular)	600-700	650 mg
	Sodium bicarbonate (granular)	300-500	367 mg
	Sodium carbonate, anhydrous	20-60	40 mg
	Flavoring agent (optional)	10-50	25 mg
	Colorant (optional)	0-10	5 mg
20	Sodium benzoate	5-15	7.5 mg
	Water	0-5	2 mg.

Premix sodium benzoate with sodium bicarbonate and alendronate sodium. Mix color with sodium carbonate. Place citric acid in a bowl of a suitable blender. Add the 2 mg water to the citric acid slowly and mix thoroughly to form a moist blend. Add to the blend, in sequence, while mixing, the sodium bicarbonate mix, and the sodium carbonate-color mix until uniformly distributed. Compress tablets using suitable size tooling. Cure the tablets, cool and package in aluminum foil.

EXAMPLE 2

	Alendronate sodium		
	(in mg alendronate acid)	5-10	5 mg or 10 mg
5	Citric acid, anhydrous (granular)	450-650	590 mg
	Sodium bicarbonate (granular)	750-1000	850 mg
	Sodium bicarbonate, powder	50-150	87 mg
	Citrus flavor (optional)	0-50	25 mg
	Water	0-25	15 mg.

10

Blend alendronate sodium, citric acid and citrus flavor in a suitable blender. Quickly add all of water and mix until a workable mass is formed. Granulate through a suitable screen using a granulator. Spread evenly on a paper-lined tray or a fluid bed dryer. Place a dried granulation in a suitable blender and add powder sodium bicarbonate. Mix well. Compress tablets using a suitable flat face beveled edge tooling. Package in aluminum foil or aluminum tubes.

20

EXAMPLE 3

	Alendronate sodium	
	(in mg alendronate acid)	10 mg
	Mannitol, direct compression grade	165 mg
25	Sodium bicarbonate	850 mg
	Citric acid	530 mg
	Sodium benzoate	15 mg

Mix alendronate sodium and sodium benzoate in a suitable mixer. Add mannitol and continue mixing till uniform. Add sequentially citric acid and sodium bicarbonate. Roller compact the powder mix followed by granulation using a suitable screen. Compress the granulation using a suitable tooling. Package the tablets in aluminum foil.

35

EXAMPLE 4

	Alendronate sodium (in mg alendronate acid)	2.5-10	10 mg
5	Citric acid, anhydrous (granular)	400-800	600 mg
	Sodium bicarbonate (granular)	1000-2000	1500 mg
	Sodium carbonate, anhydrous	20-80	40 mg
	Flavoring agent (optional)	0-50	25 mg
	Colorant (optional)	0-15	5 mg
10	Sodium benzoate	25-175	150 mg
	Xylitol	100-300	200 mg
	Sweetner (aspartame)	1-10	2.5 mg

Part 1. Blend in a suitable mixer xylitol, alendronate and
15 400 mg sodium bicarbonate. Roller compact this mixture.

Part 2. Blend in a suitable mixer sodium benzoate, citric
acid, 900 mg sodium bicarbonate, color, flavor and sweetener; then roller
compact this mixture.

Part 3. Granulate Parts 1 and 2 through a suitable screen,
20 add remaining powders and mix.

Compress the granulation using a suitable tooling. Package the tablets in
aluminum foil.

WHAT IS CLAIMED IS:

1. An effervescent pharmaceutical formulation comprising as an active ingredient a bisphosphonate selected from the groups consisting of: alendronate, cimadronate, clodronate, EB-1053, etidronate, ibandronate, neridronate, olpadronate, pamidronate, risedronate, tiludronate, YH 529, zoledronate, pharmaceutically acceptable salts and esters of the foregoing, and mixtures of the foregoing;
and
an acid source selected from the group consisting of: citric acid, tartaric acid, malic acid, fumaric acid, adipic acid, succinic acid; an anhydride of said acids; an acid salt selected from the group consisting of sodium dihydrogen phosphate, disodium dihydrogen pyrophosphate and sodium acid sulfite; and mixtures of the acids, anhydrides and acid salts; a carbonate source selected from the group consisting of: sodium bicarbonate, sodium carbonate, potassium bicarbonate, potassium carbonate, sodium sesquicarbonate, sodium glycine carbonate, and mixtures thereof;
a binder selected from the group consisting of a non-reducing sugar, polyvinyl pyrrolidone, sodium chloride, sodium benzoate and sodium sulfate;
a lubricant selected from the group consist of stearic acid salts, powdered sodium benzoate, L-leucine, sodium laurel sulfate and PEG; and
optionally, one or more additional agents selected from the group consisting of flavoring agents, colorants, and sweeteners.
2. An effervescent pharmaceutical formulation according to Claim 1 wherein the active ingredient comprises alendronate or a pharmaceutically acceptable salt thereof as an active ingredient and further comprising:

an acid source selected from the group consisting of: citric acid, tartaric acid, malic acid, fumaric acid, adipic acid, succinic acid; an anhydride of said acids; an acid salt selected from the group consisting of sodium dihydrogen phosphate, disodium dihydrogen pyrophosphate and sodium acid sulfite; and mixtures of the acids, anhydrides and acid salts; a
5 carbonate source selected from the group consisting of: sodium bicarbonate, sodium carbonate, potassium bicarbonate, potassium carbonate, sodium sesquicarbonate, sodium glycine carbonate, and mixtures thereof;

10 wherein the acid source is present in an equal or excess amount compared to the carbonate source, based on molecular equivalent amounts;

a binder selected from the group consisting of mannitol, sorbitol, polyvinyl pyrrolidone, lactose, sodium chloride, sodium
15 benzoate and sodium sulfate;

a lubricant selected from the group consist of stearic acid salts, powdered sodium benzoate, L-leucine, sodium laurel sulfate and PEG; and

20 optionally, one or more additional agents selected from the group consisting of flavoring agents, colorants, and sweeteners.

3. A formulation according to Claim 2 wherein the amount of alendronate or salt thereof is from 1-80 mg.

25 4. A formulation according to Claim 3 wherein the acid source is citric acid, tartaric acid or mixtures thereof.

5. A formulation according to Claim 3 wherein the carbonate source is sodium bicarbonate, sodium carbonate, and mixtures
30 thereof.

6. A formulation according to Claim 5 wherein the mole ratio of citric acid to bicarbonate is from 1:1 to 1:3.

7. A formulation according to Claim 5 wherein the binder is mannitol, lactose, dextrose or sorbitol.

8. A formulation comprising (in mg):

5	Alendronate sodium	
	(in mg alendronic acid)	5-10 mg
	citric acid	600-750
	sodium bicarbonate	300-500
	sodium carbonate	20-60
10	sodium benzoate	5-15
	water	0-5.

9. A formulation according to Claim 8, further comprising a flavoring agent or a colorant.

15

10. A formulation comprising (in mg):

	Alendronate sodium	
	(in mg alendronic acid)	5-10
	citric acid	450-650
20	sodium bicarbonate (total)	800-1150.

11. A formulation comprising (in mg):

	Alendronate sodium	
	(in mg alendronic acid)	2.5-10
25	citric acid	400-800
	sodium bicarbonate	1000-2000
	sodium carbonate	20-80
	sodium benzoate	25-175
	xylitol	100-300.

30

12. A formulation made by the process of combining:

	Alendronate sodium	
	(in mg alendronic acid)	2.5-10
	citric acid	400-800
5	sodium bicarbonate	1000-2000
	sodium carbonate	20-80
	sodium benzoate	25-175
	xylitol	100-300.